

The Hilbert space of Chern-Simons theory on the cylinder. A Loop Quantum Gravity approach

Clisthenis P. Constantinidis^{*,1}, Gabriel Luchini^{*,2}
and Olivier Piguet^{*,1}

March 2, 2010

¹Departamento de Física, Universidade Federal do Espírito Santo (UFES)
Vitória, ES, Brazil

²Instituto de Física de São Carlos (IFSC), Universidade de São Paulo (USP)
São Carlos, SP, Brasil

E-mails: cpconstantinidis@pq.cnpq.br, gabriel.luchini@ursa.ifsc.usp.br,
opiguet@pq.cnpq.br

Abstract

As a laboratory for loop quantum gravity, we consider the canonical quantization of the three-dimensional Chern-Simons theory on a noncompact space with the topology of a cylinder. Working within the loop quantization formalism, we define at the quantum level the constraints appearing in the canonical approach and completely solve them, thus constructing a gauge and diffeomorphism invariant physical Hilbert space for the theory. This space turns out to be infinite dimensional, but separable.

1 Introduction

Chern-Simons (CS) topological theory [1, 2, 3, 4, 5] is one of the simplest field theoretic systems showing general covariance, i.e. full invariance under space-time

^{*}Work supported in part by the Conselho Nacional de Desenvolvimento Científico e Tecnológico – CNPq (Brazil) and by the PRONEX project No. 35885149/2006 from FAPES – CNPq (Brazil).

diffeomorphisms, which characterizes it as a background independent theory. Far simpler than genuine gravitation theory in higher dimensional space-times and already very well studied [3, 6, 7, 8, 9, 10], this 3-dimensional topological theory however deserves a study in the “loop quantization” framework introduced for the canonical quantization of General Relativity and described in the books and review papers [11, 12, 13, 14, 15], where references to the numerous original papers can be found. As far as the authors know, the application of this scheme to the CS theory has not yet appeared in the literature, although it has been successfully applied to various other topological models of the Schwarz type [1, 5], such as BF theories [16], and to low dimensional gravitation theories [17, 18, 19] – which are also Schwarz topological theories.

The starting point is the canonical quantization program of Dirac [20, 21] which, in the case of a generally covariant theory, involves a Hamiltonian purely made of constraints. One first constructs a kinematical Hilbert space where the fields are represented by operators, and then selects the physical states as the vectors satisfying the constraints. Since the constraint operators generate the gauge invariances of the theory, the physical states are in fact the gauge invariant vectors.

A peculiarity of CS theory is that the space components of the gauge connection A form a pair of conjugate variables, so that the wave functional Ψ in the Schrödinger picture is a function of one of these components, let us say A_1 . Then A_2 is represented by a functional derivative. On the other hand, in topological theories of the Schwarz type, diffeomorphism invariance is a simple consequence of gauge invariance [4], at least at the classical level. One would therefore expect that, applying the Gauss constraint which ensures spatial gauge invariance, one would automatically ensure invariance under spatial diffeomorphisms and thus determine the physical Hilbert space. We will however see that, due to the necessity of choosing a polarization, i.e. choosing which component of A plays the role of a coordinate and which one plays the role of a momentum, diffeomorphism invariance is not automatic and must be implemented at the end as another constraint.

The canonical formalism requires the space-time topology to be that of $\mathbb{R} \times \Sigma$ where $\mathbb{R}$ stands for the time dimension and Σ for a space slice. In order to proceed with some details a topology for space must also be specified. We will choose that of a noncompact space, namely of a cylinder: $\Sigma = \mathbb{R} \times S^1$. As we will see, this choice leads to an infinite dimensional physical Hilbert space. To the best of our knowledge, nonperturbative quantization in the case of a noncompact space slice has not yet been considered in the literature, except the case of $\mathbb{R}^2$, which leads to a 1-dimensional Hilbert space [8, 9]. Apart of the latters, published results¹ concern compact closed spaces, where Hilbert space is finite dimensional, as well as spaces with punctures or with boundary, where exist the local degrees of freedom of a two-dimensional conformal theory [3, 7].

¹See quoted references, and [5] for more references.

Let us remind that the approach of the present paper is of the type “quantize first and then apply the constraints” and has been applied to CS theory by various authors, in particular by [8, 9]. It has to be contrasted with the approach “reduce first the classical phase space by imposing the constraints there, and then quantize”, which has been used in particular by Witten in his pioneering paper [3] (see also [10]). Both approaches may lead to inequivalent quantum field theories [8].

We will essentially follow the reference [8] for the statement of the problem and will use some of its results and notations. Our own contribution is an explicit construction of the kinematical and physical Hilbert spaces with a well defined internal product. We will restrict ourselves to a compact semi-simple Lie group of gauge invariance, typically $SU(2)$, in order to avoid the difficulties which may arise in the noncompact case in the definition of the internal product [22].

After briefly recalling in Section 2 some basic facts on the classical CS theory in the canonical framework, we proceed to the construction of the quantum state space in Section 3. Concluding remarks are presented in Section 4.

2 Classical Chern-Simons theory *à la* Dirac

Chern-Simons theory, being a fully constrained theory, may conveniently be treated using Dirac’s method [20, 21, 23], which in turn, arises from the canonical point of view. In this Section devoted to the classical theory, we follow² the reference [8]. The action is given by

$$S = -\kappa \int_{\mathcal{M}} d^3x \epsilon^{\mu\nu\rho} \text{Tr} \left(A_\mu \partial_\nu A_\rho + \frac{2}{3} A_\mu A_\nu A_\rho \right). \quad (2.1)$$

The dynamical field $A = A_\mu^I \tau_I dx^\mu$ with $\mu = 0, 1, 2$, is a Lie algebra valued connection 1-form. The gauge Lie group G will be assumed to be compact and semi-simple. The generators of the Lie algebra³ satisfy the product $[\tau_I, \tau_J] = f_{IJ}^K \tau_K$, where f_{IJ}^K are the algebra’s structure constants and $\text{Tr}(\tau_I \tau_J) = -\frac{1}{2} \delta_{IJ}$. The tensor f_{IJK} is completely antisymmetric in its three indices. The κ appearing in the above equation is the coupling constant which is well-known to be quantized [2, 3], $\kappa = \frac{n}{4\pi}$, due the gauge invariance of the quantum path integral.

The model is a generally covariant theory, integration in (2.1) being performed on a “space-time” 3-manifold $\mathcal{M}$ without metric structure. Hence there is no *a priori* notion of “time”. However, in a canonical approach, a time variable is introduced through the hypothesis that space-time has the topological structure of $\Sigma \times \mathbb{R}$, where

²The canonical formalism for Chern-Simons theory may be found in [24].

³In the case of the $SU(2)$ group which will be used throughout this paper, one has $\tau_I = -\frac{i}{2} \sigma_I$ ($I=1,2,3$), and $f_{IJK} = \epsilon_{IJK}$.

“space” is given by the 2-dimensional hypersurface Σ and “time” by the real line $\mathbb{R}$. The action then reads

$$S = -\kappa \int_{\mathbb{R}} \int_{\Sigma} dt d^2x \epsilon^{ab} \text{Tr} \left(\dot{A}_a A_b + A_0 F_{ab} \right), \quad (2.2)$$

with the spatial curvature given by $F_{ab}^I = \partial_a A_b^I - \partial_b A_a^I + f_{JK}^I A_a^J A_b^K$, $a, b, \dots = 1, 2$.

An analysis according to the Dirac-Bergman’s algorithm [20, 21] leads to a symplectic structure corresponding to the following Dirac brackets:

$$\{A_a^I(x), A_b^J(y)\} = \frac{1}{\kappa} \epsilon_{ab} \delta^{IJ} \delta^2(x - y), \quad (2.3)$$

and to the Hamiltonian

$$H = G(\varepsilon) := \int_{\Sigma} d^2x \varepsilon^I(x) G^I(x) \approx 0, \quad (2.4)$$

where ε^I is an arbitrary test function and G^I the Gauss constraint

$$G^I := -\frac{\kappa}{2} \epsilon^{ab} F_{ab}^I \approx 0. \quad (2.5)$$

We see that this is a completely constrained system and that the space components A_1^I and A_2^I form a pair of conjugate variables. The Gauss constraint is first class and its Dirac bracket algebra reproduces the Lie algebra of the gauge group:

$$\{G(\varepsilon), G(\varepsilon')\} = G(\varepsilon \times \varepsilon'), \quad (\varepsilon \times \varepsilon')^I := f_{JK}^I \varepsilon^J \varepsilon'^K. \quad (2.6)$$

It generates the space gauge transformations

$$\{A_a^I(x), G(\varepsilon^I)\} = D_a \varepsilon^I(x) = \partial_a \varepsilon^I(x) + f_{JK}^I A_a^J(x) \varepsilon^K(x). \quad (2.7)$$

We finally remember that diffeomorphism invariance – which is explicitly verified by the original action (2.1), can be shown to follow directly from gauge invariance. In particular, implementation of the Gauss constraint guaranties invariance under spatial diffeomorphisms. Indeed, such a diffeomorphism is given in the infinitesimal form by the Lie derivative $\mathcal{L}_{\xi}$ along a spatial vector field $\xi = (\xi^z, \xi^{\theta})$, and one easily checks that it is equal to a gauge transformation with parameter $\xi^a A_a^I$, up to a term proportional to the Gauss constraint:

$$\mathcal{L}_{\xi} A_a = \xi^b F_{ba} + D_a(\xi^b A_b) \approx D_a(\xi^b A_b). \quad (2.8)$$

3 Construction of the Hilbert space

Quantization *à la* Dirac is performed in two steps, namely construct first a kinematical Hilbert space $\mathcal{H}_{\text{kin}}$ based on the phase space coordinates provided by the gauge connection, more precisely its space components A_a , and then select the physical states through the constraints which, in the present case, are given by (2.5).

3.1 The kinematical Hilbert space

Working in the Schrödinger picture, we choose the spatial components $A_\theta^I := A_1^I$ as the generalized coordinates and $A_z^I := A_2^I$ as the generalized momenta obeying, as operators, the canonical commutation relations

$$[\hat{A}_\theta^I(x), \hat{A}_z^J(y)] = \frac{i}{\kappa} \delta^{IJ} \delta^2(x - y) \quad (3.1)$$

corresponding to the Dirac bracket relations (2.3) of the classical theory⁴. The indices z and θ refer to a special choice of coordinates $z \in \mathbb{R}$ and θ ($0 \leq \theta \leq 2\pi$) adapted to the topological configuration we are demanding for the 2-dimensional space Σ , namely that of a cylinder $\mathbb{R} \times S^1$. States are described by wave functionals – in Dirac's notation: $\Psi[A_\theta] = \langle A_\theta | \Psi \rangle$. The field operators act on these states as

$$\langle A_\theta | \hat{A}_\theta^I(x) | \Psi \rangle = A_\theta^I(x) \Psi[A_\theta], \quad \langle A_\theta | \hat{A}_z^I(x) | \Psi \rangle = \frac{1}{i\kappa} \frac{\delta}{\delta A_\theta^I(x)} \Psi[A_\theta]. \quad (3.2)$$

This choice of polarization, namely of A_θ^I as the configuration variables and of A_z^I as the momentum variables will turn out to be the more adequate to our aim of describing quantum states as gauge invariant functions of holonomies.

Up to now everything we have written remains purely formal until we define an integration measure in configuration space allowing us to define an internal product between the state vectors $|\Psi\rangle$. Before doing this, let us already examine the effect of the Gauss constraint at this formal level. It acts on the wave functionals as the operator

$$\hat{G}^I \Psi[A_\theta] = i \left(\frac{\partial}{\partial \theta} \frac{\delta}{\delta A_\theta^I} + f_{IJ}^K A_\theta^J \frac{\delta}{\delta A_\theta^K} \right) \Psi[A_\theta] + \kappa \left(\frac{\partial}{\partial z} A_\theta^I \right) \Psi[A_\theta]. \quad (3.3)$$

A particular solution of the Gauss constraint

$$\hat{G}^I \Psi[A_\theta] = 0, \quad (3.4)$$

is provided by the phase [8]

$$\Psi_\circ[A_\theta] = e^{2\pi i \alpha_\circ}, \quad (3.5)$$

with

$$\alpha_\circ = 4\pi\kappa \int_{\tilde{\Sigma}} d^3x \, w(g) - \frac{\kappa}{2\pi} \int_{\Sigma} d^2x \, \text{Tr}(A_\theta g^{-1} \partial_z g), \quad (3.6)$$

where the first integral is performed on a 3-dimensional manifold $\tilde{\Sigma}$ whose boundary is space Σ , and $g \in G$ is defined in terms of A_θ by $A_\theta = g^{-1} \partial_\theta g$. The integral of w over $\tilde{\Sigma}$,

$$\int_{\tilde{\Sigma}} d^3x \, w := \frac{1}{24\pi^2} \int_{\tilde{\Sigma}} d^3x \, \left(\epsilon^{\mu\nu\rho} \text{Tr}(g^{-1} \partial_\mu g \, g^{-1} \partial_\nu g \, g^{-1} \partial_\rho g) \right), \quad (3.7)$$

⁴We take $\hbar = 1$.

called the Wess-Zumino-Witten action, is an integer for the case where G is a non-abelian compact group, and consequently, a singlevalued wave functional requires then the quantization of the coupling constant:

$$\kappa = \frac{n}{4\pi}, \quad n \in \mathbb{Z}. \quad (3.8)$$

Moreover [8], the general solution of (3.4) is given by

$$\Psi[A_\theta] = \Psi_\circ[A_\theta] \psi^{\text{inv}}[A_\theta], \quad (3.9)$$

where $\psi^{\text{inv}}[A_\theta]$ – the “reduced functional” – is “ θ -gauge invariant”, i.e., obeys the condition

$$i \left(\frac{\partial}{\partial \theta} \frac{\delta}{\delta A_\theta^I} + f_{IJ}{}^K A_\theta^J \frac{\delta}{\delta A_\theta^K} \right) \psi^{\text{inv}}[A_\theta] = 0, \quad (3.10)$$

where the functional derivative operator in the left-hand side is the first part of the Gauss constraint operator (3.3) and generates the θ -gauge transformation

$$\delta A_\theta^I = D_\theta \varepsilon^I, \quad \delta A_z^I = f_{JK}{}^I A_z^J \varepsilon^K. \quad (3.11)$$

Before going to the construction of the reduced functionals ψ^{inv} in (3.9), let us proceed to the definition of the kinematical Hilbert space $\mathcal{H}_{\text{kin}}$ and of the field operators acting in it. In order to have a Hilbert space we need a well defined scalar product,

$$\langle \Psi_1 | \Psi_2 \rangle := \int \mathcal{D}\mathcal{A} \, \overline{\Psi}_1[A] \Psi_2[A], \quad (3.12)$$

with an integration measure $\mathcal{D}\mathcal{A}$ in the space of the connections defined in such a way that the scalar product is compatible with the Gauss constraint partially solved by (3.9). We therefore write

$$\Psi[A_\theta] = \Psi_\circ[A_\theta] \psi[A_\theta], \quad (3.13)$$

and will look for $\psi[A_\theta]$. The Gauss constraint takes the form (3.10) in terms of $\psi[A_\theta]$, i.e., it expresses the invariance of the latter under the θ -gauge transformations (3.11), which will be implemented in Subsection 3.3.

In the spirit of LQG, we change the focus from the Lie algebra-valued connection A_θ , which transforms inhomogeneously under gauge transformations accordingly to (3.11), to the holonomies of A_θ , which are elements of the gauge group. An holonomy $h(\gamma_{z,\theta_1,\theta_2})$ is defined over some constant z path $\gamma = [\theta_1, \theta_2]$ in Σ as

$$h(\gamma_{z,\theta_1,\theta_2}) = \mathcal{P} e^{\int_{\theta_1}^{\theta_2} d\theta \, A_\theta^I(\theta,z) \tau^I}, \quad (3.14)$$

where $\mathcal{P}$ stands for “path ordered product”. This choice is motivated by the fact that under a gauge transformation of A_θ , this holonomy transforms homogeneously:

$$h(\gamma_{z,\theta_1,\theta_2}) \longmapsto g^{-1}(z, \theta_2) \, h(\gamma_{z,\theta_1,\theta_2}) \, g(z, \theta_1) \quad (3.15)$$

We have to restrict to constant z paths because any other paths would involve the component A_z , which does not enter as an argument of the wave functional.

Let us now define the space Cyl consisting of all the wave functionals of the form (3.13) with ψ given by arbitrary finite linear combinations of complex valued functions f of the holonomies:

$$\psi[A_\theta] = \psi_{\Gamma,f}[A_\theta] = f(h(\gamma_{z_1,\theta_1,\theta'_1}), \dots, h(\gamma_{z_k,\theta_k,\theta'_k}), \dots, h(\gamma_{z_N,\theta_N,\theta'_N})), \quad (3.16)$$

where Γ denotes the “graph” defined as the (finite) set of paths

$$\Gamma = \{\gamma_{z_k,\theta_k,\theta'_k}, \quad k = 1, \dots, N\}. \quad (3.17)$$

(See Figure 3.1.) Elements of Cyl are called “cylindrical functions”. A cylindrical function is thus a functional of A_θ which, apart from the phase factor $\Psi_\circ$, depends only on the values that its argument takes on the graph Γ .




Figure 1: The construction of graphs in the space Σ , with the topology of a cylinder. The figure at left shows a single curve at some constant value for the coordinate z , and the figure at right is the representation of a particular graph in Σ , made from the disjoint union of curves at different “heights” z_k .

Since we are dealing with wave functionals of the form (3.13) with now $\psi = \psi_{\Gamma,f}$ being a function of a finite number of holonomies, which in turn are elements of the gauge group, we can now count on the invariant Haar measure of the gauge group G to define a scalar product in Cyl. Let us first define it for two state vectors⁵ $|\Gamma, f\rangle$ and $|\Gamma, f'\rangle$ associated with the same graph Γ (3.17), f and f' being two arbitrary integrable functions on $G^{\times N}$:

$$\langle \Gamma, f | \Gamma, f' \rangle = \int \prod_{k=1}^N dh_k \overline{f(h_1 \cdots h_N)} f'(h_1 \cdots h_N), \quad (3.18)$$

where dh_k is the Haar measure used to integrate over the group element h_k . The Haar measure being normalizable for a compact group, the scalar product is well

⁵We use Dirac’s notation, where the wave functional may be written as $\Psi_{\Gamma,f}[A_\theta] = \langle A_\theta | \Gamma, f \rangle$.

defined, provided the functions f on the group defining the cylindrical functions as in (3.16), are integrable. We shall take a normalized measure: $\int dh(h) = 1$.

In the general case where the two vectors are associated to two distinct graphs Γ and Γ' , we can still apply the definition (3.18), but for the graph $\Gamma'' = \Gamma \cup \Gamma'$. Indeed, the functions $f(h_1 \cdots h_N)$ and $f'(h_1 \cdots h_{N'})$ may be considered both as functions of the N'' arguments corresponding to the paths of Γ'' .

We note that this scalar product is a genuine scalar product for the vector space Cyl – whose elements are the wave functionals (3.13) with $\psi[A_\theta]$ given by (3.16). Indeed, it is a positive definite sesquilinear form in Cyl .

Finally, the kinematical Hilbert space $\mathcal{H}_{\text{kin}}$ is the Cauchy completion $\overline{\text{Cyl}}$ of Cyl with respect to the scalar product we have just defined.

Remarks

- In contrast to theories usually considered in the loop quantization framework, such as BF theories, gravity, etc., the wave functionals which are solutions of the Gauss constraint do not depend here only on the holonomies of the connection. Indeed, the phase factor $\Psi_\circ$ in (3.13), being an integral of A_θ -dependent group elements over all space, cannot be expressed in terms of holonomies. However this does not cause any problem since our definition of the scalar product is independent of this phase factor.
- By construction, the scalar product is gauge invariant and it is also invariant under the diffeomorphisms which preserve the polarization choice.

3.2 A basis for the kinematical Hilbert space

From the Peter-Weyl theorem [25], the wave functional corresponding to a graph Γ and to a function f according to (3.16) can be expanded in terms of a group representation basis as⁶:

$$\begin{aligned} \Psi_{\Gamma,f}[A_\theta] &= \Psi_\circ[A_\theta] f(h(\gamma_{z_1,\theta_1,\theta'_1}), \dots, h(\gamma_{z_k,\theta_k,\theta'_k}), \dots, h(\gamma_{z_N,\theta_N,\theta'_N})) \\ &= \Psi_\circ[A_\theta] \sum_{\vec{j}, \vec{\alpha}, \vec{\beta}} c_{\beta_1 \cdots \beta_N, j_1 \cdots j_N}^{\alpha_1 \cdots \alpha_N} R_{\alpha_1}^{j_1, \beta_1}(h(\gamma_{z_1,\theta_1,\theta'_1})) \cdots R_{\alpha_N}^{j_N, \beta_N}(h(\gamma_{z_N,\theta_N,\theta'_N})), \end{aligned} \quad (3.19)$$

where $R_\alpha^{j,\beta}(h)$ is a matrix element of the spin j unitary irreducible representation of the group element h . Thus, to every path $\gamma_{z_k,\theta_k,\theta'_k}$ of the graph Γ we associate a spin j_k representation of $G = \text{SU}(2)$.

⁶From now on we restrict ourselves to the gauge group $\text{SU}(2)$.

Moreover, the terms in the right-hand-side form an orthonormal system of vectors $|\Gamma, \vec{j}, \vec{\alpha}, \vec{\beta}\rangle$:

$$\begin{aligned} & \langle \Gamma, \vec{j}, \vec{\alpha}, \vec{\beta} | \Gamma, \vec{j}', \vec{\alpha}', \vec{\beta}' \rangle \\ &= \int \prod_{k=1}^N dh_k \overline{R_{\alpha_1}^{j_1, \beta_1}(h_1) \cdots R_{\alpha_N}^{j_N, \beta_N}(h_N)} R_{\alpha'_1}^{j'_1, \beta'_1}(h_1) \cdots R_{\alpha'_N}^{j'_N, \beta'_N}(h_N) = \delta_{\vec{j}\vec{j}'} \delta_{\vec{\alpha}\vec{\alpha}'} \delta_{\vec{\beta}\vec{\beta}'} . \end{aligned} \quad (3.20)$$

The contribution corresponding to a graph with a spin 0 path is equal to the contribution corresponding to the smaller graph obtained by erasing this path. In order to avoid a double counting, we will only include in the summation (3.19), from now on, terms where all the j_k are different from zero. In view of this restriction, it is clear that the set of vectors

$$\left\{ |\Gamma, \vec{j}, \vec{\alpha}, \vec{\beta}\rangle ; \forall \Gamma, \forall (\vec{j}, \vec{\alpha}, \vec{\beta}) \right\} \cup \left\{ |\emptyset\rangle \right\} , \quad (3.21)$$

where the vector $|\emptyset\rangle$ is associated to the empty graph $\Gamma = \emptyset$, corresponding to the wave functional $\Psi_\emptyset[A_\theta] = \langle A_\theta | \emptyset \rangle = 1$, is an orthonormal basis of the kinematical Hilbert space $\mathcal{H}_{\text{kin}}$. As a consequence of the Peter-Weyl formula (3.20), valid for any spin including spin 0, vectors associated to different graphs are orthogonal. Thus the kinematical Hilbert space is a direct sum:

$$\mathcal{H}_{\text{kin}} = \bigoplus_{\Gamma} \mathcal{H}_{\Gamma} , \quad (3.22)$$

where $\mathcal{H}_{\Gamma}$ is the Hilbert space associated with the graph Γ , and the summation is made over all possible graphs. Whereas each $\mathcal{H}_{\Gamma}$ is a separable Hilbert space, this is obviously not the case for $\mathcal{H}_{\text{kin}}$.

3.3 Solution of the Gauss constraint

In the preceding section we saw that the change of the configuration variable from A_θ to the holonomies $h[A_\Gamma]$ allows for a well defined scalar product and consequently, a Hilbert space. Although one cannot define a local operator $\hat{A}_\theta$ acting in $\mathcal{H}_{\text{kin}}$ because of the discontinuity of the scalar product [11, 12, 13], one can do it for the holonomy operator $\hat{h}[A_\theta, \gamma]$ associated to a path γ at constant z :

$$\langle A_\theta | \hat{h}[A_\theta, \gamma] | \Gamma, f \rangle = h[A_\theta, \gamma] \langle A_\theta | \Gamma, f \rangle , \quad (3.23)$$

for any basis vector $|\Gamma, f\rangle$. The resulting right-hand-side is indeed an element $|\tilde{\Gamma}, \tilde{f}\rangle$ of Cyl , associated to a new graph $\tilde{\Gamma}$ equal to the union $\Gamma \cup \{\gamma\}$ and to the function $\tilde{f} = \hat{h}[A_\theta, \gamma]f$.

Although the Gauss constraint – the infinitesimal generator of gauge transformations – expressed by (3.10), exists as a well defined operator⁷, we will impose the constraint in the form of the invariance under all finite gauge transformations as in the standard LQG approach [11, 12, 13, 14]. More precisely, taking into account the phase factor $\Psi_\circ$ in (3.13) or (3.19), we will demand the gauge invariance of the reduced wave functional ψ given by (3.16) in terms of holonomies. In view of the transformation law (3.15), it is clear that the gauge invariant reduced functionals are functions of the trace of the holonomies along closed paths (cycles), i.e. of the Wilson loops

$$h_z = \text{Tr } h(\gamma_{z,\theta,\theta}) , \quad (3.24)$$

which depend on the coordinate z , but not on the base angle θ . Thus the graphs are now sets C of cycles, each cycle being characterized by its “height” z : $C \leftrightarrow (z_1, \dots, z_n)$.

This condition of gauge invariance defines the Hilbert space $\mathcal{H}_{\text{Gauss}}$. Its basis is the orthonormal set of “spin network” vectors $|C, J\rangle$ which are given by the traces of the basis vectors of $\mathcal{H}_{\text{kin}}$:

$$\langle A_\theta | C, J \rangle = \Psi_\circ(A_\theta) \prod_{k=1}^n \chi^{j_k}(h_{z_k}) , \quad \text{with } \chi^j(h_z) = \text{Tr } R^j(h_z) , \quad (3.25)$$

where J stands for $(j_1, \dots, j_n)$ and R^j is the spin j representation of $\text{SU}(2)$. One has the orthonormality property

$$\langle C, J | C', J' \rangle = \delta_{C,C'} \delta_{J,J'} . \quad (3.26)$$

Let us define S_0 as the vector space of all finite linear combinations of spin-networks,

$$|\Psi\rangle = \sum_{n=1}^N c_n |C_n, J_n\rangle . \quad (3.27)$$

The Hilbert space $\mathcal{H}_{\text{Gauss}}$ is the Cauchy completion of S_0 . It decomposes in orthogonal subspaces in a way analogous to $\mathcal{H}_{\text{kin}}$:

$$\mathcal{H}_{\text{Gauss}} = \bigoplus_C \mathcal{H}_{\text{Gauss}}^C . \quad (3.28)$$

The Hilbert space $\mathcal{H}_{\text{Gauss}}^C$ associated to a single graph C is separable, but $\mathcal{H}_{\text{Gauss}}$ is not, since the graphs are indexed by n -arrays of real numbers.

3.4 Diffeomorphism invariance

We observe that $\mathcal{H}_{\text{Gauss}}$ is not invariant under the space diffeomorphisms. In particular, its basis vectors $|C, J\rangle$, depend explicitly on the z coordinates of the cycles

⁷The proof follows the one given by [12, 13, 14] in a more general context.

constituting the graph C and are therefore not invariant under changes of the coordinate z . (They are however invariant under changes of the coordinate θ .) This has to be contrasted with the situation in the classical theory, where the fulfilment of the Gauss constraint automatically ensures full space diffeomorphism invariance (see (2.8)). We note from (2.8) that a diffeomorphism along the z coordinate generated by a vector field $\xi = (\xi^z, 0)$, acts on the configuration variable A_θ as $\mathcal{L}_\xi A_\theta \approx D_\theta(\xi^z A_z)$, i.e. as a gauge transformation with parameter $\xi^z A_z$. However, when applied to a wave functional, A_z must be replaced by the operator defined in (3.2). Such a “gauge transformation” was not contemplated when we solved the Gauss constraint in Subsection 3.3. Therefore, we have still to implement this part of diffeomorphism invariance, namely invariance under the z -diffeomorphisms

$$z' = z'(z), \quad \theta' = \theta. \quad (3.29)$$

The more general diffeomorphisms generated by vectors $\xi = (\xi^z, \xi^\theta)$ – which modify the polarization – are left aside for the time being.

In the same way as we have proceeded with the Gauss constraint, we will impose invariance under the *finite* z -diffeomorphisms (3.29) due to the difficulty of defining their infinitesimal generator.

We will follow the standard group averaging method [11, 12, 13], based here on the Gel’fand triple [26] $S_0 \subset \mathcal{H}_{\text{Gauss}} \subset S'_0$, where S_0 is the subspace of finite linear combinations of spin-networks defined at the end of the preceding subsection, dense in $\mathcal{H}_{\text{Gauss}}$, and S'_0 its dual, whose elements are the complex valued linear functionals Φ of S_0 :

$$\Phi : S_0 \rightarrow \mathbb{C}, \quad \Psi \mapsto \langle \Phi, \Psi \rangle \in \mathbb{C}, \quad (3.30)$$

where we use Schwartz notation $\langle \cdot, \cdot \rangle$ for functionals [27]. The scalar product (3.18) in $\mathcal{H}_{\text{Gauss}}$ being explicitly invariant under all space diffeomorphisms, any such diffeomorphism ϕ is represented by a unitary operator U_ϕ . The action of ϕ in S'_0 is then defined by duality:

$$\langle U_\phi \Phi, \Psi \rangle = \langle \Phi, U_{\phi^{-1}} \Psi \rangle. \quad (3.31)$$

We will concentrate on the z -diffeomorphisms (3.29).

The z -diffeomorphism invariant states are now given by vectors of S'_0 constructed from any vector $|\Psi\rangle$ of S_0 by applying to it the operator P_{Diff} – a functional “projector”:

$$P_{\text{Diff}} : S_0 \rightarrow S'_0, \quad \langle P_{\text{Diff}} \Psi, \Psi' \rangle = \sum_{\Psi''} \langle \Psi'' | \Psi' \rangle, \quad \forall |\Psi'\rangle \in S_0, \quad (3.32)$$

where the sum is done over all the vectors $|\Psi''\rangle$ of S_0 which may be obtained from $|\Psi\rangle$ by a z -diffeomorphism: $|\Psi''\rangle = U_\phi |\Psi\rangle$. The sum in (3.32) is always finite. Indeed, the vectors $|\Psi\rangle$ and $|\Psi''\rangle$ are both finite superpositions of spin-networks vectors – see (3.27):

$$|\Psi\rangle = \sum_{m=1}^M c_m |C_m, J_m\rangle, \quad |\Psi''\rangle = \sum_{n=1}^N c''_n |C''_n, J''_n\rangle. \quad (3.33)$$

Since two spin-network vectors are orthogonal if their respective graphs are different, the summation in (3.32) is restricted to those $|\Psi''\rangle$ which have at least one graph C_n'' in its expansion which coincides with a graph C_m of the expansion of $|\Psi\rangle$. But, in the present theory where space is one-dimensional, given a spin-network vector $|C_m, J_m\rangle$, there are only two such spin-network vectors: $|C_m, J_m\rangle$ itself and the vector $|C_n'', J_n''\rangle$, with C_n'' , obtained from the first one by inverting the order of the “edges” of the graph C_m , i.e. of the coordinates z_k which label these edges – with the restriction that the spins j_k'' attributed to the edges still match after this inversion.

The functionals $P_{\text{Diff}}\Psi$ defined in this way are z -diffeomorphism invariant, as a consequence of the equation (3.31) which defines the way elements of S'_0 transform. They span by definition the vector space $\mathcal{H}_{\text{Phys}} \subset S'_0$, which we will show to be the physical Hilbert space of the theory.

We observe that the only element of this space which is also an element of $\mathcal{H}_{\text{Gauss}}$ is the trivial state $|0\rangle$ – the “vacuum” – defined by

$$\langle A_\theta | 0 \rangle = \Psi_\circ(A_\theta), \quad (3.34)$$

where $\Psi_\circ$ is the phase factor (3.5), which is obviously diffeomorphism invariant, in particular under the z -diffeomorphisms considered here.

We define now the scalar product in $\mathcal{H}_{\text{Phys}}$ by

$$\langle P_{\text{Diff}}\Psi_1 | P_{\text{Diff}}\Psi_2 \rangle := \langle P_{\text{Diff}}\Psi_1, \Psi_2 \rangle \quad (3.35)$$

where $|\Psi_1\rangle, |\Psi_2\rangle \in S_0$. This product is independent of the particular state $|\Psi_2\rangle$ we used to define $|P_{\text{Diff}}\Psi_2\rangle$.

By construction (see (3.32)), vectors of $\mathcal{H}_{\text{phys}}$ only depend on the equivalence classes of vectors of S_0 under z -diffeomorphisms. In particular, a vector defined by (3.32) from a spin-network $|C, J\rangle$ does not depend on the particular positions z_k of the cycles constituting the graph C , but only on the number of such cycles – and of the spin values associated to each of them. Following the LQG terminology [11], let us call such a vector an s-knot and denote it by $|j_1, \dots, j_N\rangle \equiv |J\rangle$:

$$|J\rangle = P_{\text{Diff}}|C, J\rangle \quad (3.36)$$

The scalar product of two s-knots is given by

$$\langle J | J' \rangle = \langle j_1, \dots, j_N | j'_1, \dots, j'_{N'} \rangle = \delta_{NN'} \left(\delta_{j_1 j'_1} \cdots \delta_{j_N j'_{N'}} + \delta_{j_N j'_1} \cdots \delta_{j_1 j'_{N'}} \right). \quad (3.37)$$

The second term in the right-hand side is due to the existence, mentioned above, of z -diffeomorphisms which preserve a graph but reverse the order of its cycles. Thus, provided one identifies a vector $|j_1, j_2, \dots, j_N\rangle$ with its “reversed” $|j_N, \dots, j_2, j_1\rangle$, the s-knot states provide an orthonormal basis of $\mathcal{H}_{\text{Phys}}$.

The vectors (3.36) being completely characterized by finite sets of half-integer numbers $J = \{j_1, j_2, \dots, j_N\}$, are clearly invariant under *all diffeomorphisms*, beyond being solutions of the Gauss constraint. In particular, they do not depend on the choice of the polarization. $\mathcal{H}_{\text{phys}}$ is thus the physical state space of the Chern-Simons theory on a cylinder. The set of s-knots vectors being countable, this Hilbert space is separable.

4 Conclusions

We found in Chern-Simons theory a great opportunity to discuss the main ideas of loop quantum gravity, concerning the method by itself. In fact, this theory fits exactly in the framework which was used to think on quantum gravity in the beginning [28], where the quantum states were generated by Wilson loops of the Ashtekar connection [29], the so called “loop states”.

What we have concretely done here is a continuation of [8]. In this reference the authors showed the path to get a physical quantum state based on the implementation of the Gauss constraint, which could not be implemented in the usual way because of the difficulty to define the quantum operators based on the choice of a Schrödinger representation. In order to really define a Hilbert space, we needed, after having fixed a particular topology (as suggested in [8]) for the spatial slice Σ , to change the configuration variables from the connection to its holonomies, following the prescriptions of Loop Quantum Gravity. The result, for the chosen topology, is a physical Hilbert space whose basis is indexed by a nonnegative integer N and, for each N , by an ordered set of N half integer numbers $j_1, \dots, j_N$ – the spins.

Our choice of A_θ – the connection component in the direction of the compact space dimension – as the configuration variable, and of A_z – the component in the noncompact direction – as the conjugate variable, induced the breaking of general covariance. This is the reason why the Gauss constraint was not sufficient to assure full diffeomorphism invariance, which was finally recovered by imposing the invariance under the diffeomorphisms along the z coordinate.

One could wonder on our choice (3.13), with (3.16), for the elements of the vector space Cyl whose Cauchy completion yields the kinematical Hilbert space $\mathcal{H}_{\text{kin}}$. The main reason for doing so is that it allows for a well defined scalar product and for a simple solution of the Gauss constraint in terms of θ -gauge invariant functions of the $z = \text{constant}$ holonomies. Another approach⁸, which would apparently be more in the spirit of loop quantum gravity, would consist in taking, as a kinematical space, the functions of the $z = \text{constant}$ holonomies, without the phase factor Ψ_o , then trying to construct the Gauss constraint operator in this space and finally taking its

⁸We thank a referee for suggesting it.

kernel. Even if manageable, this way would certainly be much more cumbersome than the one we have followed.

We have taken $SU(2)$ as the gauge group. Generalization to other compact Lie groups looks straightforward. The consideration of more general topologies, as well as the construction of physical observables along the same lines are left for future works.

Acknowledgments We thank Laurent Freidel, Camillo Imbimbo and Galen Sotkov for most valuable informations and discussions. This work was supported in part by the Conselho Nacional de Desenvolvimento Científico e Tecnológico – CNPq (Brazil) and by the PRONEX project No. 35885149/2006 from FAPES – CNPq (Brazil).

References

References

- [1] A.S. Schwarz, *Lett. Math. Phys.* 2 (78) 247; *Commun. Math. Phys.* 67 (79) 1.
- [2] S. Deser, R. Jackiw and S. Templeton, *Phys. Rev. Lett.* 48 (1982) 975; *Annals Phys.* 140 (1982) 372; *Erratum-ibid.* 185 (1988) 406; *Annals Phys.* 281 (2000) 409.
- [3] E.Witten “Quantum Field Theory and the Jones Polynomial, *Commun. Math. Phys.* 121 (1989) 351.
- [4] E. Witten; “2 + 1 dimensional Gravity as an Exactly Soluble system”, *Nucl. Phys.* B311 (1988) 46.
- [5] D. Birmingham, M. Blau, M. Rakowski and G.T. Thompson, “Topological Field Theory”, *Phys. Rep.* 209 (1991) 129.
- [6] S. Elitzur, G. Moore, A. Schwimmer and N. Seiberg, “Remarks on the Canonical Quantization of the Chern-Simons-Witten Theory”, *Nucl. Phys.* B326 (1989) 108.
- [7] G. Moore and N. Seiberg, “Taming the Conformal Zoo”, *Phys. Lett.* B220 (1989) 422.
- [8] G.V. Dunne, R. Jackiw, C.A. Trugenberger, “Chern-Simons Theory in the Schrödinger Representation”, *Annals Phys.* 194 (1989) 197.
- [9] E.Guadagnini, M.Martellini, M.Mintchev, “Braids and Quantum Group Symmetry in Chern-Simons Theory, *Nucl. Phys.* B336 (1990) 581.

- [10] M. Bos and V.P. Nair, “Coherent State Quantization of Chern-Simons Theory”, *Int. J. Mod. Phys. A*5 (1990) 959.
- [11] C. Rovelli; “Quantum Gravity”, Cambridge University Press, 2004 [arXiv:gr-qc/9710008].
- [12] T. Thiemann, “Modern Canonical Quantum General Relativity”, Cambridge Monographs on Mathematical Physics, 2007.
- [13] A. Ashtekar and J. Lewandowski, “Background Independent Quantum Gravity: A Status Report”, *Class. Quantum Grav.* 21 (2004) R53, [arXiv:gr-qc/0404018].
- [14] M. Han, W. Huang and Y. Ma “Fundamental structure of loop quantum gravity”, *Int. J. Mod. Phys. D*16 (2007) 1397, [arXiv:gr-qc/0509064].
- [15] A. Perez, “Introduction to loop quantum gravity and spin foams”, Lectures given at 2nd International Conference on Fundamental Interactions, Domingos Martins, Espirito Santo, Brazil, 6-12 Jun 2004, published in “Proceedings of Second International Conference on Fundamental Interactions 2004”, p. 221, [arXiv:gr-qc/0409061].
- [16] J.C. Baez, “An Introduction to spin foam models of quantum gravity and BF theory”, *Lect. Notes Phys.*543:25-94,2000, [arxiv:gr-qc/9905087].
- [17] K. Noui and A. Perez, “Three-dimensional loop quantum gravity: Physical scalar product and spin foam models”, *Class. Quantum Grav.* 22 (2005) 1739, [arXiv:gr-qc/0402110].
- [18] J. Manuel Garcia-Islas, “Observables in three-dimensional quantum gravity and topological invariants”, *Class. Quantum Grav.* 21 (2004) 3933, [arXiv:gr-qc/0401093].
- [19] C.P. Constantinidis, A. Perez and O. Piguet, “Quantization of the Jackiw-Teitelboim model”, *Phys. Rev. D*79 (2009) 084007, [arXiv:0812.0577[gr-qc]].
- [20] P.A.M. Dirac, “Lectures on Quantum Mechanics”, Belfer Graduate School of Science, Yeshiva University Press, New York, 1964;
P.A.M. Dirac, “Lectures on Quantum Mechanics”, Belfer Graduate School of Science, Dover, Mineola, NY, 2001.
- [21] M. Henneaux, C. Teitelboim; “Quantization of Gauge Systems”, Princeton University Press, 1994.
- [22] L. Freidel and E. Livine, “Spin networks for noncompact groups”, *J. Math. Phys.* 44 (2003) 1322, [arXiv:hep-th/0205268].
- [23] H.J. Matschull “Dirac’s canonical quantization program”, [arXiv:quant-ph/9606031].

- [24] S. Deser and X. Xiang, *Phys. Lett.* B263 (1991) 39.
- [25] A.O. Barut, “Theory of Group Representations and Applications”, Polish Scientific Publishers, Warszawa, 1977.
- [26] I. M. Gel’fand and N. J. Vilenkin, “Generalized Functions, vol. 4: Some Applications of Harmonic Analysis. Rigged Hilbert Spaces”, Academic Press, New York, 1964;
R. de la Madrid, “The role of the rigged Hilbert space in Quantum Mechanics”, [arXiv:quant-ph/0502053v1].
- [27] Laurent Schwartz, “Théorie des distributions”, Hermann, 1997.
- [28] C. Rovelli and L. Smolin, “Knot Theory and Quantum Gravity”, *Phys. Rev. Lett.* 61 (1988) 1155;
C. Rovelli and L. Smolin, “Loop Space Representation of Quantum General Relativity”, *Nucl. Phys.* B331 (1990) 80.
- [29] A. Ashtekar, “New Hamiltonian Formulation of General Relativity”, *Phys. Rev.* D36 (1987) 1587.